

# CELLOBIOHYDROLASE HYDROLYZES CRYSTALLINE CELLULOSE ON HYDROPHOBIC FACES\*

Yu-San Liu<sup>1,2</sup>, John O. Baker<sup>1</sup>, Yining Zeng<sup>1,2</sup>, Michael E. Himmel<sup>1,2</sup>, Thomas Haas<sup>1,2</sup>, and Shi-You Ding<sup>1,2</sup>

<sup>1</sup>Biosciences Center, National Renewable Energy Laboratory, 1617 Cole Boulevard, Golden, Colorado 80401

<sup>2</sup>BioEnergy Science Center, Oak Ridge National Laboratory, P.O. Box 2008, Oak Ridge, Tennessee 37831

Running head: CBH I hydrolyzes cellulose on hydrophobic faces

Address correspondence to: Shi-You Ding, Ph.D., Biosciences Center, National Renewable Energy Laboratory, 1617 Cole Boulevard, Golden, CO 80401

303-384-7758. Fax: 303-384-7752; E-mail: [Shi.you.Ding@nrel.gov](mailto:Shi.you.Ding@nrel.gov)

**Biodegradation of plant biomass is a slow process in nature, and hydrolysis of cellulose is also widely considered to be a rate-limiting step in the proposed industrial process of converting lignocellulosic materials to biofuels. It is generally known that a team of enzymes including endo- and exo-cellulases as well as cellobiases are required to act synergistically to hydrolyze cellulose to glucose. The detailed molecular mechanisms of these enzymes have yet to be convincingly elucidated. In this report, Atomic Force Microscopy (AFM) is used to image in real-time the structural changes in *Valonia* cellulose crystals acted upon by the exo-cellulase cellobiohydrolase I (CBH I) from *Trichoderma reesei*. Under AFM, single enzyme molecules could be observed binding only to one face of the cellulose crystal, apparently the hydrophobic face. The surface roughness of cellulose began increasing after adding CBH I, and the overall size of cellulose crystals decreased during an eleven-hr period. Interestingly, this size reduction apparently occurred only in the width of the crystal, whereas the height remained relatively constant. In addition, the measured cross-section shape of cellulose crystal changed from asymmetric to nearly symmetric. These observed changes brought about by CBH I action may constitute the first direct visualization supporting the idea that the exo-cellulase selectively hydrolyzes the hydrophobic faces of cellulose. The limited accessibility of the hydrophobic faces in native cellulose may contribute significantly to the rate-limiting slowness of cellulose hydrolysis.**

Natural cellulose is a bundle of linear  $\beta$ -1,4-linked glucan chains held tightly in a crystalline structure by the cumulative effect of many inter- and intra-chain hydrogen bonds. Cellulose produced by higher plants is the most abundant biopolymer on Earth, accounting for 40-60% by dry-weight of plant cell walls. In addition to the traditional uses of cellulose in the paper, food, and textile industries, the new concept of biofuels produced from lignocellulosic biomass is considered a promising route to sustainable energy production. Unfortunately, lignocellulosic material is intrinsically recalcitrant to chemical and enzymatic breakdown to simple sugars that can be fermented to liquid fuels. A deeper understanding of biomass recalcitrance will be required for the potential of lignocellulosic biofuels to be realized (1).

In native plant cell walls, cellulose exists as nanometer-scale microfibril networks embedded in matrices of other biopolymers such as hemicelluloses, pectins and lignins. Chemical pretreatment processes are often required to remove or relocate these 'other' matrix polymers, thereby exposing the cellulose to a follow-up enzymatic hydrolysis to produce glucose. The cellulases are a class of enzymes, produced mainly by cellulolytic fungi and bacteria, that catalyze hydrolysis of the  $\beta$ -1,4-glucosidic bonds that link the glucosyl units of cellulose. Cellulases have been classified on the basis of their modes of action on the substrate into three distinct classes that react synergistically: (i) the endo- $\beta$ -(1,4)-glucanases that cleave the cellulose chain at internal positions to produce new chain ends (2), (ii) the exo- $\beta$ -(1,4)-D-glucanases that cleave successive cellobiosyl units from the ends of cellulose and (iii) the  $\beta$ -D-glucosidases that

hydrolyze cellobiose to glucose. Among these types of cellulases, the exo-glucanases appear to catalyze most of the bond-cleavages in the saccharification of crystalline cellulose, and are usually the major component of cellulase preparations, especially in the case of current fungus-derived commercial enzymes.

Cellulose substrates isolated from different sources, though all composed of linear  $\beta$ -1,4-linked glucose polymers, differ significantly in structure as measured in terms of amorphous content, crystallinity, and size and shape of crystallites (3); these structural differences may substantially affect susceptibility to cellulase action. In addition, depending on variations of the patterns of inter- and intra-chain hydrogen-bonds, cellulose may form different crystalline allomorphs (e.g., cellulose I $_{\alpha}$ , I $_{\beta}$ , II, III and IV), upon which the binding and activities of cellulases may be different. Endo-glucanases acting alone may attack the non-crystalline regions effectively, but effective attack on the crystalline portion of cellulose requires in addition the synergistic action of the exo-glucanases. The fact that cellulose structure affects cellulase activities has long been documented (4), but experimental data presented in the literature have been subject to considerable uncertainty in interpretation of the relationships between cellulose structure and cellulase activities due to disagreements in identification and quantification of cellulose structure by different analytical methods (5). In the current study, cellulose crystals from the cell wall of a green alga, *Valonia ventricosa*, were intensively characterized by means of atomic force microscopy (AFM). Single, continuously-monitored crystals of the characterized material were used as substrates in assays visualizing and quantifying the effects of catalysis by *Trichoderma reesei* CBH I (Cel7A) on the structure of cellulose crystals. The ultimate objective of this study is to elucidate the reaction mechanism of cellulase on true crystalline cellulose.

The cellulase system produced by the filamentous fungus *T. reesei* is widely used as a commercial enzyme mixture in industry. This system contains two exo-glucanases -- cellobiohydrolases CBH I and CBH II, which belong to glycoside hydrolase families 7 and 6 respectively, along with five endo-glucanases (EG

I-V). It is generally believed that hydrolysis of crystalline cellulose to cellobiose is the rate-limiting step of cellulose degradation, which requires at least two type of synergistic reactions, endo-exo cooperation between endoglucanases (EGs) and cellobiohydrolases (CBHs), and exo-exo cooperation between CBH I and II, which remove cellobiosyl residues from, respectively, the reducing and non-reducing ends of cellulose chains (6). Among the enzymes present in the cellulase preparation from *T. reesei*, CBH I is the most abundant component, accounting for more than 60% of the total protein (7). CBH I is structurally modular, consisting of a family 7 glycoside hydrolase catalytic module and a family 1 carbohydrate-binding module (*TrCBM1*), connected by a proline- and threonine-rich, highly glycosylated linker peptide. In a proposed (8) mechanism for the action of CBH I on crystalline cellulose, the surface-binding family 1 CBM first binds to the planar surface of cellulose. The reducing end of one cellulose chain is then threaded into the active-site tunnel of the catalytic module. The cellulose chain is advanced "processively" through the tunnel, two glucosyl residues at a time, as catalytic residues in the tunnel catalyze the hydrolysis of every second  $\beta$ -1,4-glucosidic bond to depolymerize the chain to free cellobiose units. This last proposed feature is based primarily on the analytical finding that CBH I produces mainly cellobiose, as well as on the structure of the catalytic module (8) and the CBM (9). A recent study employing fast-AFM of CBH I on a cellulose crystal appears supportive of this proposed mechanism (10).

Over the last decade, researchers have investigated cellulase activity and cellulose-cellulase interaction through a combination of biochemical methods and wet chemistry, as well as imaging techniques such as fluorescence microscopy (11-14), transmission electron microscopy (TEM) (15,16) and scanning electron microscopy (SEM) (17). AFM has also been used to characterize the effects of different types of cellulases upon cellulose structure (10,13,18-20). Previously we have studied the interactions with cellulose of both complete cellulase enzymes (13) and their binding modules (14,21), by employing Single-Molecule Spectroscopy (SMS) techniques, such as Total Internal Reflection Fluorescence Microscopy and AFM. It has been demonstrated

that AFM is a powerful analytical tool that can be used to obtain highly accurate images of the cellulose surface under physiological conditions with nanometer resolution (22,23). The sample can be imaged without physical (freezing) or chemical (fixation) treatments; the images thus obtained therefore reflect “native” structures. However, up until now, limited information has been obtained by real-time observation of biological events in time frames long enough to capture significant effects of cellulase action. In the present study, we use AFM to image in real-time the morphological changes occurring in single crystals of *Valonia* cellulose as a result of hydrolysis by *T. reesei* CBH I. The *Valonia* cellulose I has mostly I<sub>a</sub> structure, and is a widely accepted native crystalline cellulose standard. Our objective was to learn how the enzyme affects the cellulose morphology at the nanometer scale.

## EXPERIMENTAL PROCEDURES

*T. reesei* CBH I purification and *Valonia* *ventricosa* cell culture - CBH I was purified from a seven-day broth of *T. reesei* grown on Sigmacell 50 (Sigma-Aldrich Corporation, St. Louis MO, USA) as carbon source. The detailed protocol of CBH I purification has been described elsewhere (24). The cell culture of the green alga *V. ventricosa* was purchased from UTEX The Culture Collection of Algae at The University of Texas at Austin, and was grown in enriched seawater medium (UTEX) at 23°C under a photo regime of 16 hr light at 80 to 100 ft-c intensity alternated with 8 hr periods in the dark. The cells were harvested by centrifugation, and were stored at -20°C.

*Preparation of Valonia cell wall cross-section for AFM* - Frozen whole *Valonia* cells were fixed and embedded using the following protocol, with all steps performed in a microwave at 30°C under vacuum except where noted. Samples were fixed in 2.5% (v/v) glutaraldehyde in phosphate buffered saline solution (PBS), washed 3X with PBS, fixed in 1% (w/v) OsO<sub>4</sub> in PBS, washed 3X with distilled H<sub>2</sub>O, and dehydrated with increasing concentrations of ethanol (30%, 60%, 90%, 3 × 100%, diluted in distilled H<sub>2</sub>O). Fixed cells were then infiltrated with EMBED 812 resin (EMS, Hatfield, PA) in increasing concentrations of resin (7%, 15%, 30%, 60%, 90%, 3 × 100% resin,

diluted in ethanol), with the last two resin exchanges lasting for 48 hr each on a rotator at room temperature. The samples were transferred to flat-bottomed BEEM-type capsules (EMS, Hatfield, PA) and the resin was polymerized by heating to 60°C for 72 hr in a nitrogen-purged vacuum oven. The samples were then sectioned to about 240 nm thickness with a Diatome diamond knife (EMS, Hatfield, PA) on a Leica EM UTC ultramicrotome (Leica, Wetzlar, Germany). Sections were collected on freshly-cleaved mica disks for AFM.

*Preparation of Valonia cellulose crystals for AFM imaging in buffer and CBH I solution* - The cellulose crystals were isolated from *Valonia* cells using the previously described method (25). Purified *Valonia* cellulose crystals were stored at -20°C. Freshly cleaved mica coated with an amine polymer, poly-L-lysine, provided a flat, smooth, and stationary substrate on which the cellulose crystals were firmly immobilized to overcome the forces applied by the AFM probe and to permit high resolution of the sample structure. Five μL of poly-L-lysine solution (0.1% w/v in water, Sigma-Aldrich, St. Louis, MO) was added to freshly cleaved mica. After 5 minutes incubation, the excess poly-L-lysine solution was removed using a spin coater (Model KW-4A, Chemat Technology, Northridge, CA) operated at 500 rpm for 30 sec followed by 4000 rpm for 30 sec. After that, 5 μL of a suspension of cellulose crystals (1 mg/mL in water) was added, followed by incubation and spin-coating as in the above procedure. The coated mica was then dried in an oven at gentle heat (45°C) overnight. AFM was operated in tapping mode in buffer and in air. We found that application of a 150-μL liquid droplet (enzyme solution or buffer) to the cellulose-coated mica provided a sample in aqueous environment that was sufficient for imaging for at least 10 hrs without significant artifacts caused by water evaporation. CBHI (1 mg/mL) was applied to the cellulose-coated mica in acetate buffer (20 mM acetate, 0.02% NaN<sub>3</sub>, 100 mM NaCl, pH 5.0): the same buffer, without CBHI, was used for buffer-control imaging.

*AFM and image analysis* - AFM measurements were conducted using a Multi-Mode™ scanning probe microscope with NanoScope V controller (Veeco, Santa Barbara, California). To ensure absolute stability, the AFM was located in a

specially designed laboratory with acoustic and vibration isolation. A customized Nikon optical microscope with deep focus (800X magnification) was used to aid in positioning of the AFM tip to the desired location. Silicon nitride probes (DNP) and sharp nitride lever (SNL) probes (Veeco, Santa Barbara, California) were used in tapping mode to image cellulose fibers in liquid, and DP18/HIRES/Al BS probes (MikroMasch, San Jose, California) were used in air. Images of cross section and roughness were analyzed using Nanoscope 7.30 software. Surface roughness was evaluated by flattening the images (second-order) and calculating mean roughness (Ra) and root mean squared roughness (Rq). Ra is the arithmetic average of the absolute values of the surface height deviations measured from the mean plane, expressed as

$$Ra = \frac{1}{N} \sum_{j=1}^N |Z_j|$$

whereas Rq is root mean square

average of height deviations taken from the mean image data plane, expressed as  $Rq = \sqrt{\frac{\sum (Z_i)^2}{N}}$ .

## RESULTS AND DISCUSSION

*Size and shape of Valonia cellulose crystals.* The green alga *V. ventricosa*, produces large cellulose fibrils. Cross-sections of *Valonia* cellulose crystallites appear square with sizes ranging from 10 to 20 nm as observed by TEM (26,27). The negative staining method used in those studies measured the crystallites as non-staining objects, and it was expected that the measured size of the crystallite could be smaller than that of the actual microfibril due to staining effects and resolution limitation. In the current study, the cross-section of the *Valonia* cell wall was imaged by AFM, with height (Figure 1A), phase (Figure 1B) and amplitude (Figure 1C) images being taken simultaneously. By analyzing these images, the contours of each microfibril could be clearly seen. The shape of each microfibril is hexagonal (Figure 1D); because the microfibrils are mostly irregular with two narrow sides, they can appear pentagonal in low-resolution images (Figure 1A, 1B, 1C). The sizes of individual microfibrils range from 10 to 50 nm, with maximum frequency near 35 nm. The isolated cellulose crystals were also imaged

(Figure 1E and 1F). Several crystals often form small bundles, and single crystals were measured in the cross-section profile of height images as being 15 to 40 nm in height with maximum frequency of 25 nm, and the length was measured as 200 nm to 2  $\mu$ m with maximum frequency of 1  $\mu$ m. The reason that the measured size of single isolated crystals was slightly smaller than that in native cell wall could be: 1) the measured cross-sections may not be perfectly perpendicular to the long axis of the microfibril, which would result in larger size measured than the actual one; 2) although we used a sharp tip (1 nm in apex) for the AFM imaging to minimize the tip broadening effect, it would be difficult to deconvolute such effects; or 3) the preparation process used to isolate the cellulose crystal could also be expected to cause surface damage and peeling. Nevertheless, the shape of *Valonia* cellulose observed is an irregular hexagon with two narrow sides. Incidentally, subunits corresponding to the 3-5 nm elementary fibril previously reported (28) were not observed in this study.

*CBH I binds to cellulose and moves.* Figure 2 shows a single cellulose crystal imaged in acetate buffer (as “zero-time control”, Figure 2A) and at 171, 179, 188, 196, 205, 214 and 222 minutes, respectively, (Figure 2B-H) after addition of CBH I. With the same scanning size and area, we observed that cellulose fibers had well-defined ridges and relatively clean surfaces in buffer. After addition of CBH I, there were some new features (right-hand-side of the cellulose) showing up on the cellulose crystal. These new features could be explained as CBH I enzymes binding to the cellulose surface. These new features were observed over a period of time (222 minutes), as shown in Figure 2 (B-H). The apparent size of these features is approximately a quarter of the AFM-measured height (25 nm) of the cellulose crystal. Based on published crystal structures for the CBH I catalytic domain (8), and on small-angle X-ray scattering structures for the intact enzyme (29), projection of a bound CBH I molecule away from the cellulose surface may be inferred to be between 5 and 10 nm. A productively-bound molecule (bound through both CBM and a catalytic module “threaded” upon a cellulose chain) will have a “projection height” closer to the smaller size, whereas an enzyme molecule bound to the cellulose only by the CBM

might have projection heights ranging up to approximately 10 nm. Therefore, we propose that the new features appearing on cellulose are most likely to be CBH I enzymes. Although the AFM time-resolution (frame rate ~1 frame/(8 min)) used in this study is not capable of demonstrating processive motion of specific, individual CBH I molecules, the observed new features were similar to those tracked in a recent high-speed AFM report (10). Figure 2 also shows that these features change relative locations on the cellulose surface over the time imaged. Because there are apparently two different sizes of features, with their relative distances changing during the observed time frames (Figure 2C and 2D), we further speculate that the two size-categories represent two different kinds of modules (i.e., catalytic and carbohydrate-binding modules), and that the observed changes in intermodule distance reflect conformational changes in the entire CBH I molecule as CBH I catalyzes bond-cleavage and moves along on a cellulose chain. Rigorous statistical analysis of continuous real-time imaging using AFM will be required to confirm this speculation.

*Cellulose surface roughness increases with CBH I reaction.* AFM is capable of measuring the surface structure at atomic resolution. We analyzed the structural changes in the cellulose surface concomitant with the action of CBH I. One of the features obtained from AFM measurements is the degree of roughness, which can be used to analyze changes in the surface brought about by friction, adhesion, and catalytic activity. There are many mathematical approaches to calculating surface roughness from AFM images, including probability height distribution, fractal analysis, mean roughness (Ra), and root-mean-squared roughness (Rq). Ra and Rq are the most commonly reported of the measures of surface roughness that can be extracted from AFM images. Figure 3 represents roughness measurement of the cellulose crystal shown in Figure 2. We found that cellulose surface roughness increased (in both Ra and Rq cases) by about 0.3 nm after addition of CBH I, which further confirms that the apparent morphological changes of cellulose observed by AFM were the results of CBH I reaction. The roughness remained almost constant during a subsequent 274 minutes of measurement in the presence of CBH I (Figure

3). *Valonia* celluloses are known to be primarily  $I_{\alpha}$  in crystalline form (30). The distance between  $I_{\alpha}$  cellulose sheets ( $d_{110}$ ) is 0.39 nm, which is similar to the measured surface roughness increases resulting from CBH I action, indicating that CBH I enzymes affect only the surface layer of cellulose crystals. Previously it has been demonstrated that the *TrCBM1* binds to the hydrophobic faces (1 1 0) of cellulose (21,31,32), and these faces are as narrow as 2-4 nm (14,32). Presumably the hydrophobic faces consist of more than one cellulose chain, thus the roughness change may indicate that the cellulose chains are hydrolyzed individually.

*Cellulose crystal size decreases during CBH I hydrolysis.* Continuous images of cellulose were taken in real-time during incubation with either buffer or CBH I solution. The size and shape of the cellulose crystal were analyzed by plotting the cross-section profiles of each image frame (Figure 4A and 4B). No change was observed during an 11-hr incubation in buffer (Figure 4A). After addition of CBH I, on the other hand, Figure 4B shows size reduction of the observed area taken from the average of two successive frames with 50 cross section lines in each frame (based on 512 x 512 scan lines), which demonstrated that the average width of the cellulose crystal decreased, but the average height remained relatively constant. The apparent reduction of cellulose specifically on one side of the crystal may imply that the CBH I tends to hydrolyze cellulose from certain surfaces. It has been previously reported that the family 1 CBM from CBH I binds only to certain faces of cellulose crystal, i.e., the (1 1 0) faces (or the hydrophobic faces) in the case of *Valonia*  $I_{\alpha}$  cellulose (28). It could be further speculated that CBH I hydrolysis also occurs only on the hydrophobic faces as discussed below.

The cross-section area of each single cellulose crystal was further integrated based on AFM image section profiles, and size reduction of 20% was estimated based on 11 hrs of measurement (Figure 4C), whereas there was no change observed for the same period of time in buffer (Figure 4A). The hydrolysis rate of CBH I could not be calculated, because 1) it is not known how many CBH I molecules are involved in the period of 11 hrs hydrolysis, and 2) it is difficult to estimate how many cellulose chains are hydrolyzed in the 20% size reduction.

*CBH I hydrolysis mechanism.* Figure 5 shows a schematic diagram of an anisotropic hexagonal cellulose crystal and predictions of scanning profiles produced during AFM imaging. In this diagram, the cellulose crystal is initially 25 nm in width and is shaped based on cellulose I<sub>α</sub> structure with two hydrophobic faces (1 1 0) both 3 nm wide. Most likely the broader hydrophilic faces will attach to the mica surface when a single cellulose crystal makes contact with the mica surface during AFM sample preparation. The schematic of the AFM tip is generated based on a 1 nm radius for the tip-apex and the general shape provided by the manufacturer. According to this proposed orientation of the crystal on the mica surface, both hydrophobic (1 1 0) and (-1 -1 0) faces, are exposed and susceptible to enzymatic hydrolysis. AFM, however, can probe only the surfaces geometrically accessible to the tip, so for the situation shown in Figure 5, only the hydrolytic removal of material from the (1 1 0) face will show up in scan profiles as a difference between scans taken before and after CBH I hydrolysis. The schematic diagram shown in Figure 5 therefore provides a quite likely explanation for the one-sided reduction of the cellulose crystal width in Figure 4B. In addition, Figure 5 illustrates a reasonable explanation for the observation that no significant changes in measured crystal height are detected during enzyme hydrolysis, contrasted with substantial changes in crystal width.

It has been documented that CBH I carries out exo-attack on cellulose from its reducing end generating sharpened tips (16). Figure 6 shows a succession of real-time AFM amplitude images (9 min per scan) of the end of the same cellulose crystal, beginning ("zero time" in figure legend) 6 hr after the addition of CBH I. The amplitude mode yields clear images with high contrast. During the course of the enzyme hydrolysis, we observed various shape changes at the end of the cellulose crystal, such as from smooth (Figure 6A, 0 min), to sawtoothed (Figure 6B, 9 min), and then to nicked (Figure 6C, 26 min). The fiber end appeared smooth again in Figure 6G (249 min), but with a sharper angle than that in Figure 6A. The observed tip sharpening is similar to that reported previously (16) where CBH I erodes bacterial cellulose and renders the reducing end pointed in TEM images. We extend the results of

the earlier work by providing real-time images of irregularities produced in cellulose-crystal tips by CBH I-catalyzed hydrolytic erosion. For the same reason illustrated in Figure 5, AFM only detects one side of sharpening, whereas TEM image can show sharpening from both sides. In this study, operating AFM in an aqueous environment allows us to image the process under physiological conditions without intervening chemical/physical treatments. The observed phenomenon is thus closer to that occurring in nature. A control experiment was done by imaging cellulose fiber ends exposed to acetate buffer only. A series of real-time AFM amplitude images were taken starting at six hrs after adding acetate buffer solution rather than CBH I solution. No observable changes were found in either short-time (0 to 34 min with ~9 min frame rate) or long-time scale (360 min) (Figure 7). The ends of isolated cellulose crystals sometimes appear to be nicked branches (Figure 7), which could have resulted from the harsh preparation conditions (strong acid/base or high temperature) of cellulose isolation.

## CONCLUSIONS

Atomic force microscopy has been used to make real-time measurements, both qualitative and quantitative, of the morphological changes in single crystals of cellulose during hydrolysis by *T. reesei* CBH I. We demonstrated structural changes including decrease in width, increases of surface roughness, and changes in shape of fiber ends. We have observed that the cross-section of *Valonia* cellulose crystal is apparently an irregular hexagon with two narrow hydrophobic faces that are binding surfaces for TrCBM1, and are likely to be the surfaces from which CBH I-catalyzed reactions remove cellulose chains. To the best of our knowledge, this is the first direct measurement of CBH I hydrolysis of crystalline cellulose that supports the hypothesis that CBH I hydrolyzes only the hydrophobic faces of cellulose. We further speculate that development of pretreatment approaches aimed at modification of cellulose to increase the accessibility of the hydrophobic planes to enzyme catalyst may lead to further improvements in the efficiency of enzymatic saccharification of biomass.

## REFERENCES

1. Himmel, M. E., Ding, S. Y., Johnson, D. K., Adney, W. S., Nimlos, M. R., Brady, J. W., and Foust, T. D. (2007) *Science* **315**, 804-807
2. Reese E.T., S. R. G. H., Levinson H.S. (1950) *J. Bacteriol.* **59**, 485-497
3. Montanari, S., Rountani, M., Heux, L., and Vignon, M. R. (2005) *Macromolecules* **38**, 1665-1671
4. Fan, L. T., Lee, Y. H., and Beardmore, D. H. (1980) *Biotechnol. Bioeng.* **22**, 177-199
5. Park, S., Baker, J. O., Himmel, M. E., Parilla, P. A., and Johnson, D. K. (2010) *Biotechnol. Biofuels* **3**, 10
6. Henrissat, B., Driguez, H., Viet, C., and Schulein, M. (1985) *Bio-Technology* **3**, 722-726
7. Enari, T. M., and Nikupaavola, M. L. (1987) *Crit. Rev. Biotechnol.* **5**, 67-87
8. Divne, C., Stahlberg, J., Reinikainen, T., Ruohonen, L., Pettersson, G., Knowles, J. K. C., Teeri, T. T., and Jones, T. A. (1994) *Science* **265**, 524-528
9. Kraulis, P. J., Clore, G. M., Nilges, M., Jones, T. A., Pettersson, G., Knowles, J., and Gronenborn, A. M. (1989) *Biochemistry* **28**, 7241-7257
10. Igarashi, K., Koivula, A., Wada, M., Kimura, S., Penttila, M., and Samejima, M. (2009) *J. Biol. Chem.* **284**, 36186-36190
11. Helbert, W., Chanzy, H., Husum, T. L., Schulein, M., and Ernst, S. (2003) *Biomacromolecules* **4**, 481-487
12. Moran-Mirabal, J. N., Santhanam, N., Corgie, S. C., Craighead, H. G., and Walker, L. P. (2008) *Biotechnol. Bioeng.* **101**, 1129-1141
13. Liu, Y. S., Luo, Y. H., Baker, J. O., Zeng, Y. N., Himmel, M. E., Smith, S. J., and Ding, S. Y. (2010) *Proc. SPIE* **7571**, 757103
14. Ding, S. Y., Xu, Q., Ali, M. K., Baker, J. O., Bayer, E. A., Barak, Y., Lamed, R., Sugiyama, J., Rumbles, G., and Himmel, M. E. (2006) *BioTechniques* **41**, 435-443
15. Bayer, E. A., Chanzy, H., Lamed, R., and Shoham, Y. (1998) *Curr. Opin. Struct. Biol.* **8**, 548-557
16. Boisset, C., Frascini, C., Schulein, M., Henrissat, B., and Chanzy, H. (2000) *Appl. Environ. Microbiol.* **66**, 1444-1452
17. Bohn, A., Fink, H. P., Ganster, J., and Pinnow, M. (2000) *Macromol. Chem. Phys.* **201**, 1913-1921
18. Lee, I., Evans, B. R., and Woodward, J. (2000) *Ultramicroscopy* **82**, 213-221
19. Liu, H., Fu, S. Y., Zhu, J. Y., Li, H., and Zhan, H. Y. (2009) *Enzyme Microb. Technol.* **45**, 274-281
20. Quirk, A., J., L., C., V., D., C., Clarke, A. J., Dutcher, J. R., and Roscoe, S. G. (2010) *Langmuir* **26**, 7
21. Liu, Y. S., Zeng, Y. N., Luo, Y. H., Xu, Q., Himmel, M. E., Smith, S. J., and Ding, S. Y. (2009) *Cellulose* **16**, 587-597
22. Baker, A. A., Helbert, W., Sugiyama, J., and Miles, M. J. (2000) *Biophys. J.* **79**, 1139-1145
23. Ding, S. Y., and Himmel, M. E. (2006) *J. Agric. Food Chem.* **54**, 597-606
24. Baker, J. O., Mitchell, D. J., Grohmann, K., and Himmel, M. E. (1991) Thermal Unfolding of Trichoderma reesei CBH I. in *Enzymes in Biomass Conversion* (Leatham, G. F., and Himmel, M. E. eds.), American Chemical Society, Washington DC. pp 313-330
25. Imai, T., Putaux, J. L., and Sugiyama, J. (2003) *Polymer* **44**, 1871-1879
26. Chanzy, H., Henrissat, B., Vuong, R., and Revol, J. F. (1986) *Holzforschung* **40**, 25-30
27. Revol, J. F. (1982) *Carb. Polymers* **2**, 123-134
28. Gardner, K. H., and Blackwell, J. (1971) *J. Polymer Science Part C: Polymer Symposia* **36**, 327-340
29. Abuja, P. M., Schmuck, M., Pilz, I., Tomme, P., Claeysens, M., and Esterbauer, H. (1988) *Eur. Biophys. J. Biophys.* **15**, 339-342

30. Sugiyama, J., Vuong, R., and Chanzy, H. (1991) *Macromolecules* **24**, 4168-4175
31. Lehtio, J., Sugiyama, J., Gustavsson, M., Fransson, L., Linder, M., and Teeri, T. T. (2003) *Proc. Natl. Acad. Sci. U. S. A.* **100**, 484-489
32. Xu, Q., Tucker, M. P., Arenkiel, P., Ai, X., Rumbles, G., Sugiyama, J., Himmel, M. E., and Ding, S. Y. (2009) *Cellulose* **16**, 19-26

## FOOTNOTES

\*This work was supported by the U.S. Department of Energy, Office of Energy Efficiency and Renewable Energy, the Office of the Biomass Program for the work to develop enzymes, and the U.S. DOE Office of Science, Office of Biological and Environmental Research through the BioEnergy Science Center (BESC), a DOE Bioenergy Research Center, for the work on Atomic Force Microscopy (AFM) visualization and analysis.

The abbreviations used are: AFM, atomic force microscopy; CBH I, cellobiohydrolase I; CBM, cellulose-binding module; EG, endoglucanase; DNP, silicon nitride probe; SNL, sharp nitride lever.

## FIGURE LEGENDS

**Fig. 1.** Atomic force micrographs of *Valonia* cell wall microfibrils. A layered-structure is shown in height (A), phase (B), and amplitude (C) images taken simultaneously in 1 x 1  $\mu\text{m}$  scan size of native *Valonia* cell wall cross-section. Microfibrils are nearly parallel-aligned, and each 5-8 layers of microfibrils form a lamella. Zoomed-in phase image (200 x 200 nm, D) shows the cross-section shape of individual microfibrils are mostly anisotropic hexagonal. Isolated *Valonia* cellulose crystals were also imaged in acetate buffer. Height images of 5 x 5  $\mu\text{m}$  scan size (E) shows small bundles formed by several crystals, and zooming-in on the selected area from (E) shows individual crystals (F), the size of which is about 25 nm based on height measurement. Clear ridges can be seen on the cellulose crystal. Scale bar = 100 nm (A-C), 20 nm (D), 1 $\mu\text{m}$  (E), and 30nm (F).

**Fig. 2.** 3-D rendering of time-lapse AFM amplitude images of an individual isolated *Valonia* cellulose crystal in acetate buffer (A) and at 171, 179, 188, 196, 205, 214, and 222 minutes after addition of CBH I enzyme (B-H). Features observed attached to one side of the cellulose crystal are believed to be CBH I enzyme molecules. Features appearing to have two different sizes may represent the two modules of CBH I (i.e., CD and CBM indicated by solid and open arrows, respectively). The location of CBH I on cellulose was shown to change, which indicates movement of the enzyme. The measured relative position of putative CDs and CBMs was also observed to change between approximately 5-10 nm, which is consistent with the modeled linker length, indicating that CBH I may be a dynamic structure during the catalytic process. Scale bar = 10 nm.

**Fig. 3.** Cellulose surface roughness increases after addition of CBH I. Time-lapse roughness analysis (based on Figure 2) of cellulose fibers in buffer (0 min, control experiment) and in CBH I solution (171, 179, 188, 196, 205, 214, 222 min). Roughness increases by 0.3 nm shortly after addition of CBH I, and then remains at about the same level over 3 hrs measurement. Trend lines are used to guide the eye. Ra: mean roughness, Rq: root-mean-squared roughness.

**Fig. 4.** Selected cross-section profiles of *Valonia* cellulose analyzed by AFM height image in acetate buffer (A) and CBH I solution (B) during 11 hrs of continuous imaging. Averages of 100 cross-section profiles (two successive frames with 50 cross-section lines in each frame) are plotted. In buffer, the cellulose profile remains unchanged. In CBH I solution, cellulose surface morphology changes and size decreases from one side of the crystal resulting cellulose in width changes while height remains

approximately the same. (C) shows percentage change of integrated areas relative to the first point of the sequence (the 6<sup>th</sup> hr after CBHI addition). Squares: cellulose in acetate buffer showing almost no changes; triangles: cellulose in CBH I solution showing decreases of about 20%. Note that the cellulose crystals measured in buffer and in CBH I solution are two independent, but identically-conducted experiments.

Fig. 5. A schematic diagram of AFM measurement of a cellulose crystal, in which the cross-section of a single 25 nm cellulose crystal is represented based on cellulose I<sub>α</sub> structure with narrow 3 nm hydrophobic faces (1 1 0). In this schematic representation the crystal attaches to the mica surface by its broader hydrophilic surface. If the CBH I hydrolyzes both (1 1 0) and (-1 -1 0) faces, the AFM can only measure the changes in the (1 1 0) face, resulting in cross-section profiles of AFM height imaging reduced in width from one side and unchanged in height (Figure 4). Tip scanning profiles in red and blue lines represent cellulose before and after hydrolysis, respectively. The tip and the cellulose are not in scale.

Fig. 6. Detailed structure of the end of a single cellulose crystal measured by AFM amplitude images. The cellulose crystal was treated in CBH I solution, with imaging begun after 6 hrs treatment (A), and continued for an additional 274 minutes. Representative images are shown at 9 (B), 26 (C), 43 (D), 86 (E), 240 (F), 249 (G), and 274 (H) additional minutes of CBH I treatment, respectively. At successive time points, the crystal ends appeared smooth (A), sawtoothed (B), and then nicked (C), indicating that CBH I erodes cellulose from the end of the crystal irregularly. Apparent overall end-sharpening (A-G) is also observed only on one side, due to the effects of AFM tip geometry illustrated in Figure 5. These observations support previously reported cellulose tip sharpening caused by CBH I and imaged by TEM. The AFM imaging provides real-time measurement and more details of morphological changes in crystal ends during CBH I hydrolysis. Scale bar = 20 nm.

Fig. 7. “Buffer only” control for the experiment in Figure 6: The cellulose crystal end shows no change in acetate buffer either in the short time scale (A-E) or in the long time scale (F). AFM amplitude images were taken (A) 0 min (B) 9 min (C) 17 min (D) 26 min (E) 34 min and (F) 360 min after the end of six hrs initial exposure to acetate buffer. Scale bar = 20 nm.

Figure 1




Figure 2




Figure 3




Figure 4






Figure 5



**Figure 6**



**Figure 7**

