

FEATURED ARTICLE

Two *Arabidopsis* proteins synthesize acetylated xylan *in vitro*

Breeanna R. Urbanowicz, Maria J. Peña*, Heather A. Moniz, Kelley W. Moremen and William S. York*

Complex Carbohydrate Research Center, University of Georgia, 315 Riverbend Road, Athens, GA 30602, USA

Received 4 June 2014; revised 18 July 2014; accepted 1 August 2014; published online 21 August 2014.

*For correspondence (e-mails mpena@ccrc.uga.edu; will@ccrc.uga.edu).

SUMMARY

Xylan is the third most abundant glycopolymer on earth after cellulose and chitin. As a major component of wood, grain and forage, this natural biopolymer has far-reaching impacts on human life. This highly acetylated cell wall polysaccharide is a vital component of the plant cell wall, which functions as a molecular scaffold, providing plants with mechanical strength and flexibility. Mutations that impair synthesis of the xylan backbone give rise to plants that fail to grow normally because of collapsed xylem cells in the vascular system. Phenotypic analysis of these mutants has implicated many proteins in xylan biosynthesis; however, the enzymes directly responsible for elongation and acetylation of the xylan backbone have not been unambiguously identified. Here we provide direct biochemical evidence that two *Arabidopsis thaliana* proteins, IRREGULAR XYLEM 10-L (IRX10-L) and ESKIMO1/TRICOME BIREFRINGENCE 29 (ESK1/TBL29), catalyze these respective processes *in vitro*. By identifying the elusive xylan synthase and establishing ESK1/TBL29 as the archetypal plant polysaccharide *O*-acetyltransferase, we have resolved two long-standing questions in plant cell wall biochemistry. These findings shed light on integral steps in the molecular pathways used by plants to synthesize a major component of the world's biomass and expand our toolkit for producing glycopolymers with valuable properties.

Keywords: plant cell wall, glycosyltransferase, *O*-acetyltransferase, xylan, xylan synthase, *Arabidopsis thaliana*.

INTRODUCTION

The survival of plants that dominate our environment and that are the ultimate source of most of our food depends on their ability to synthesize xylan-rich cell walls, which are required for the efficient transport of water from the soil to the leaves. Xylans are a major component of the cereals and vegetables we consume, and constitute up to one-third of the woody biomass that we use to build our homes. As a major component of the plant cell walls in both monocots and dicots, these agronomically important natural biopolymers play a central role in human health and well-being. Xylans are being increasingly used as a renewable resource for the manufacture of valuable products such as biocompatible materials (Simkovic, 2013).

Like most polysaccharides, xylans are synthesized by glycosyltransferases (GTs) that use the chemical energy in donor molecules, such as nucleotide sugars, to drive the attachment of monosaccharides to the growing polymer (Figure 1a; Lairson *et al.*, 2008). All xylan molecules produced by vascular plants have the same backbone, composed of 1,4-linked β -D-xylopyranosyl (Xyl) residues; however, the structures that decorate this backbone vary

depending on the plant tissue and species (York and O'Neill, 2008). Several enzymes that diversify xylan structure by functionally modifying its backbone have been identified. For example, glucuronic acid (GlcA) substituents are added to glucuronoxylan in dicots such as *Arabidopsis thaliana* by members of the carbohydrate active enzyme (CAZy) family GT8 (Lairson *et al.*, 2008) α -glucuronosyltransferases encoded by *GLCA SUBSTITUTION OF XYLAN (GUX)* genes (Rennie *et al.*, 2012). We have recently shown (Urbanowicz *et al.*, 2012) that the GlcA residues are further modified by GLUCURONOXILAN METHYLTRANSFERASE 1 (GXMT1) to form 4-O-methyl glucuronic acid (MeGlcA). In dicots, the most abundant substituents of the backbone Xyl residues are *O*-acetates (Teleman *et al.*, 2000). For example, the glucuronoxylan extracted from poplar wood and *A. thaliana* stems (Teleman *et al.*, 2000; Yuan *et al.*, 2013) contains approximately six times as many *O*-acetyl substituents as (Me) GlcA substituents. Recently, analysis of *A. thaliana* mutants with reduced xylan acetylation phenotypes implicated members of two protein families, REDUCED WALL




Figure 1. IRX10-L catalyzed the transfer of xylosyl residues from UDP-Xyl to a chemically labeled acceptor, Xyl₆-2AB.

(a) *In vitro* xylosyltransferase reaction scheme.

(b) MALDI-TOF MS of the products after a 16-h reaction in the absence and presence of IRX10-L. The series of annotated [M + H]⁺ ions are the result of structures with a mass difference of 132 Da, consistent with the sequential addition of xylosyl residues to the Xyl₆-2AB acceptor to generate oligosaccharides with degrees of polymerization (DP) ranging from seven to 13.

(c) 1D ¹H NMR analysis of products generated by incubating unlabeled xylohexaose for 120 h with IRX10-L and UDP-Xyl. The increased signal intensities for internal β-Xyl (shaded) relative to those of terminal and reducing β-Xyl establishes that IRX10-L catalyzes the addition of β-(1,4)-linked Xyl residues to the chain.

(d) Kinetics of xylosyl transfer to xylohexaose (Xyl₆), as determined by measuring the quantities of uridine triphosphate (UDP) formed upon transfer of the xylosyl residue to increasing quantities of Xyl₆ (0–4 mM) in the presence of IRX10-L (15.8 pmol). Three technical replicates are shown. Kinetic constants *K_m* (mM) and *V_{max}* (pmol UDP min⁻¹) were calculated by fitting the initial velocities (*V*, pmol UDP min⁻¹) as a function of the acceptor substrate concentration to the Michaelis–Menten equation using nonlinear curve fitting.

(e) The effect of divalent metals on IRX10-L activity, determined by RP-HPLC analysis. Data are averages of two biological replicates. Error bars are SDs.

ACETYLATION (RWA) and TRICHOME BIREFRINGENCE-LIKE (TBL), in xylan acetylation (Gille and Pauly, 2012; Rennie and Scheller, 2014). Phenotypic evidence suggests that RWA proteins are involved in the early steps of the acetylation process, whereas members of the TBL family are specific polysaccharide *O*-acetyltransferases (Gille and Pauly, 2012; Rennie and Scheller, 2014); however, it has not been possible thus far to obtain direct biochemical evidence showing that any of the enzymes required for *O*-acetylation of xylan *in vivo* actually catalyze the acetylation of the polysaccharide (Gille and Pauly, 2012).

It has been hypothesized (Richmond and Somerville, 2000; Sandhu *et al.*, 2009) that the enzymes responsible for the synthesis of all plant β-1,4-glycans, including xylan, are members of CAZy family GT2 (Lairson *et al.*, 2008). As a result of their structural homology with cellulose synthase (CESA; Pear *et al.*, 1996), all of these proteins are classified as cellulose synthase-like (CSL) GTs (Richmond and Somer-

ville, 2000). Enzymes in this family have been shown to catalyze β-1,4-glycan backbone synthesis for almost all other hemicellulosic polysaccharides, including mannan (CSLA; Dhugga *et al.*, 2004), xyloglucan (CSLC; Cocuron *et al.*, 2007) and 1,3;1,4-β-D-glucan (CSLF; Burton *et al.*, 2006; CSLH; Doblin *et al.*, 2009). Interestingly, the β-1,4-galactosyltransferase responsible for galactan synthesis was recently identified as a GT92 enzyme, demonstrating that CSL proteins are not always responsible for β-1,4-glycan synthesis in plants (Liwanag *et al.*, 2012). Although cell wall galactans are β-1,4-glycans, they are not configurationally homologous with cellulose, mannan, xyloglucan and xylan because the glycosidic oxygens in the main chain have an axial orientation relative to C-4 of the β-galactosyl residue. This has profound effects on the overall shape and physical properties of the polysaccharide, much the same way that a change from the β-anomeric configuration (equatorial glycosidic oxygen) to the α-anomeric configuration (axial

glycosidic oxygen) causes cellulose and starch to have completely different biophysical properties. The identification of the GT92 galactan synthase thus requires a modest refinement of the above hypothesis, as follows: 'The enzymes responsible for the synthesis of all plant β -1,4 glycans with configurational homology to cellulose are members of CAZy family GT2'.

In contrast to this hypothesis, most of the enzymes so far implicated in xylan backbone synthesis are classified as GT43 or GT47 enzymes. The genes encoding these proteins were originally discerned by phenotypic analysis of *A. thaliana* *IRREGULAR XYLEM (IRX)* mutants that produce xylan with reduced chain length (Brown *et al.*, 2005, 2009; Peña *et al.*, 2007; Wu *et al.*, 2009; Scheller and Ulvskov, 2010). Three distinct complementation pairs were identified: IRX9/IRX9-L (GT43), IRX14/IRX14-L (GT43), and IRX10/IRX10-L (GT47) (Wu *et al.*, 2009, 2010; Jensen *et al.*, 2013; Rennie and Scheller, 2014). The first gene of each pair (*IRX9*, *IRX10* and *IRX14*) is the most highly expressed in secondary cell walls. Complementation with either gene in a pair is able to rescue double mutants in which both genes have been inactivated, indicating that each gene pair encodes two isoenzymes that have very similar or identical catalytic functions (Brown *et al.*, 2009; Wu *et al.*, 2009, 2010). Xylan with a reduced chain length has been isolated from the *irx9 irx9-L* double mutant. In contrast, no xylan was detected in the stem tissues of *irx14 irx14-L* or *irx10 irx10-L* (Wu *et al.*, 2009, 2010), indicating that IRX10/IRX10-L and IRX14/IRX14-L play vital roles in the formation of this polymer. Based on the increased xylosyltransferase activity of microsomes from tobacco BY2 cells expressing both IRX9 and IRX14, Lee *et al.* (2012) proposed that these two GT43 enzymes are xylosyltransferases; however, no unambiguous biochemical evidence that establishes the precise molecular functions of any of these proteins in xylan backbone synthesis has been published. Recently, Jensen and colleagues performed transcriptional profiling of the heteroxylan-rich psyllium (*Plantago ovata* Forsk) seed mucilaginous layer, and showed that a transcript corresponding to a homolog of Arabidopsis *IRX10-L* displayed the highest expression levels of all the putative glycosyltransferases surveyed (Jensen *et al.*, 2013). Transcripts of GT43 (*IRX9* and *IRX14*) genes were found in very low abundance in the mucilage, however, but were present at levels comparable with *IRX10* homologs in the stem tissue (Jensen *et al.*, 2011, 2013), indicating that IRX10/10-L may be sufficient for xylan backbone synthesis in select tissues.

RESULTS

To determine whether IRX10-L from *A. thaliana* is capable of elongating the β -(1,4)-xylan backbone, we expressed a recombinant tagged form of this enzyme in human embryonic kidney (HEK) 293 cells and purified it from the culture medium (Figure S1a). The catalytic activity of the

enzyme was evaluated by means of a xylosyltransferase assay using uridine 5'-diphosphate-xylose (UDP-Xyl) as a sugar donor and 2-aminobenzamide β -(1,4)-xylohexaose (Xyl₆-2AB) as a fluorescently labeled acceptor (Figure 1a). Matrix-assisted laser desorption ionization time-of-flight mass spectrometry (MALDI-TOF MS) analysis of the reaction products indicated that recombinant IRX10-L catalyzes the transfer of several xylosyl residues from UDP-Xyl to Xyl₆-2AB (Figure 1b). This transfer cannot occur at the reducing end of the substrate, which is blocked by the fluorescent tag (2AB). Treatment of the synthesized polymer with a β -1,4-endoxyranase depolymerized the products (Figure S1b), consistent with the conclusion that IRX10-L catalyzes the addition of β -1,4-linked xylosyl residues to the xylan oligosaccharides. Having established a robust xylosyltransferase assay, we expressed and purified IRX9 and IRX14, as described above, and also evaluated their ability to transfer Xyl to Xyl₆-2AB; however, these proteins showed no discernible activity, alone or in combination, under these assay conditions (Figure S2). In order to obtain more rigorous characterization of the products of the reaction catalyzed by IRX10-L, we used MALDI-TOF MS (Figure S3a) and nuclear magnetic resonance (NMR) spectroscopy to fully characterize the products of a scaled-up reaction using unlabeled β -(1,4)-xylohexaose as the acceptor. The NMR spectra of these products (Figures 1c, S3b and S4a,b) showed an increase in the relative intensity of signals assigned to internal β -(1,4)-linked xylose, compared with those assigned to terminal β -Xyl and reducing α - and β -Xyl. This is consistent with the fact that all plant GT47 glycosyltransferases characterized to date employ an inverting catalytic mechanism (Geshi *et al.*, 2010), which in the case of IRX10-L transfers xylose from an α -linked donor (UDP-Xyl) to form a β -linked product (β -1,4-xylan). These data unambiguously establish that IRX10-L is a polymerizing GT that forms β -(1,4)-linkages by transferring multiple xylosyl residues from UDP-Xyl to the non-reducing end of xylooligosaccharide acceptors.

Having established that IRX10-L catalyzes xylan backbone synthesis *in vitro*, we next examined the biochemical properties of this enzyme using a bioluminescent assay to detect and quantify xylosyltransferase activity. This approach measures uridine diphosphate (UDP), which is released as a product of the transfer of Xyl from UDP-Xyl onto the elongating chain (Figure 1a). Xylosyltransferase activity was shown to be enzyme concentration-dependent in the presence of UDP-Xyl as a donor and Xyl₆ as an acceptor, with no observable release of UDP when IRX10-L was reacted with UDP-Xyl alone (Figure S1c). Furthermore, we performed experiments using various acceptors, including xylose, β -(1,4)-xylooligosaccharides with degrees of polymerization (DP) ranging from two to six and cellopentaose, a β -(1,4)-glucooligosaccharide. The data indicated that IRX10-L requires an exogenously added

acceptor with at least two sugar residues, and does not use cellopentaose as an acceptor (Figure S1d). The Michaelis constant (K_m) for IRX10-L activity using Xyl₆ as an acceptor was determined to be 1.17 mM (Figure 1d), which is similar to that reported for crude wheat microsomal membrane preparations using xylopentaose (Xyl₅) as an acceptor (2.9 mM; Kuroyama and Tsumuraya, 2001). Using the MALDI-TOF MS-based assay, we also observed that activity was optimal at pH 6.7–6.8 (Figure S5), also consistent with previous studies using microsomal membranes (Kuroyama and Tsumuraya, 2001). We tested whether IRX10-L can use other UDP sugars, including UDP-arabinopyranose, UDP-galactose, UDP-galacturonic acid, UDP-glucose and UDP-glucuronic acid, as sugar donors. No glycosyltransferase activity was detected with any of these nucleotide sugars, except for very low activity when UDP-arabinopyranose was added. We attribute this activity to the presence of UDP-Xyl, which is a known contaminant (approximately 0.35%) of the UDP-arabinopyranose preparation (CarboSource Services, personal communication).

The GTs that use nucleotide sugar donor substrates adopt two main types of three-dimensional (3D) folds. GT-A enzymes are typically divalent metal-dependent and often have a cation-binding Asp-X-Asp (DXD) signature motif, whereas most GT-B proteins are metal ion-independent (Lairson *et al.*, 2008). Although no 3D structures are yet available to place GT47 enzymes such as IRX10-L into one of the GT superfamilies, they have been predicted to adopt a GT-B-type fold (Liu and Mushegian, 2003; Lairson *et al.*, 2008). This is consistent with our data showing that purified IRX10-L is a nucleotide sugar-dependent GT that retains full catalytic activity after ion chelation with EDTA (Figure 1e). The reported increase in xylan synthase activity in wheat microsomes observed when metal ions are added in the absence of exogenous acceptors (Porchia and Scheller, 2000) suggests that other factors that contribute to xylan synthesis *in vivo* (Rennie and Scheller, 2014) may be metal-dependent.

Polymerizing GTs catalyze glycosyl chain elongation by either a distributive mechanism, where the acceptor is released after the addition of a single residue, or a processive mechanism, where the acceptor is released after the addition of several residues (May *et al.*, 2009). A distributive mechanism generates products corresponding to the addition of one or two residues during the initial stages of the reaction, followed by a Poisson distribution of larger products as the reaction progresses. In contrast, a processive mode of action produces larger oligosaccharides during the initial stages of the reaction. To distinguish between these two possibilities, we analyzed and quantified the reaction progress by reversed-phase high-performance liquid chromatography (RP-HPLC) (Figure 2a). These analyses demonstrate that IRX10-L catalyzes the




Figure 2. Characteristics of recombinant IRX10-L.

(a) RP-HPLC of the products formed by incubating IRX10-L ($0.25 \mu\text{g} \mu\text{l}^{-1}$) with UDP-Xyl (2 mM), Xyl₆-2AB (0.25 mM) and MgCl₂ (1 mM) in HEPES-HCl (75 mM), pH 6.8, for the indicated times.

(b) The time-dependent abundance distributions of oligosaccharide products generated as described in (a) and quantified by the integration of peaks in the RP-HPLC elution profiles. Data are averages of three reactions with purified protein from two biological replicates. Observed product distributions are shown for 24 h (blue) and 48 h (red) as fractional populations (dotted lines with data-point symbols); theoretical abundances (solid lines) are Poisson probabilities calculated as described in Experimental procedures.

time-dependent polymerization of xylan to produce large xylooligosaccharides after 48 h (Figure 2a). The remarkable agreement between the observed and theoretical abundances is consistent with a distributive catalytic mechanism of action for the enzyme (Figure 2b). Taken together, our data indicate that IRX10-L, unaided by other enzymes, uses a distributive mechanism to catalyze xylan synthesis *in vitro* by the repeated incorporation of xylosyl residues to the non-reducing end of the acceptor.

O-Acetyl groups are the most abundant backbone substituents of xylan in dicotyledonous plants. Despite the high degree of acetyl substitution in xylan and many other plant glycopolymers, very little is known about the process of polysaccharide *O*-acetylation in plants (Gille and Pauly, 2012). Current data are consistent with the notion that


plant polysaccharide *O*-acetylation may occur through a mechanism similar to that proposed for the *O*-acetylation of peptidoglycan by Gram-negative bacteria. For example, phenotypic evidence suggests that *Neisseria gonorrhoeae* employs a two-component system where an acetyl moiety is first translocated from a cytoplasmic acetyl donor, assumed to be acetyl-CoA, to the periplasm by an integral membrane protein, peptidoglycan *O*-acetyltransferase (PatA). *O*-acetylated PatA is proposed to then act as the acetyl donor for PatB *in vivo*, which carries out the transacetylation reaction (Moynihan and Clarke, 2010). Recently, in-depth biochemical analysis of PatB has shown that the enzyme can use several simple acyl donors *in vitro*, indicating a great deal of promiscuity in terms of donor use (Moynihan and Clarke, 2013, 2014). In plants, the integral membrane proteins from the RWA family are predicted to function in the acetylation pathway prior to TBL proteins. Recent chemotypic analysis of mutants lacking one or more RWA proteins show a global reduction in acetylation of several cell-wall polysaccharides with varying effects on individual polymers (Manabe *et al.*, 2013). These data suggest that they may function either as acetyl translocating proteins, similar to the mechanism proposed for PatA, as Golgi acetyl-CoA transporters, or in some other capacity such as components of a larger protein complex. In cases where in-depth chemical analysis has been performed on hemicellulosic polysaccharides from TBL mutants, the phenotypes are manifested as reduced acetylation in one or two specific glycopolymers (Gille *et al.*, 2011; Xiong *et al.*, 2013; Yuan *et al.*, 2013). Accordingly, phenotypic analysis

has revealed that mutating a member of the plant-specific TBL gene family (Gille and Pauly, 2012) *TRICHOME BIREFRINGENCE-LIKE 29/ESKIMO 1* (*TBL29/ESK1*) results in plants with a chemotype distinguished by xylan and mannan with a reduced degree of acetylation, indicating that TBL29 is a xylan or mannan *O*-acetyltransferase (Xiong *et al.*, 2013; Yuan *et al.*, 2013). No biochemical activity data have been published for any plant polysaccharide *O*-acetyltransferase however, and the identity of the acetyl donor and the ability of this enzyme to directly acetylate xylan or mannan have not yet been established.

To determine whether TBL29/ESK1 is capable of functioning as an *O*-acetyltransferase, we expressed and purified a recombinant tagged form of the enzyme (Figure S6a), and evaluated its capacity to transfer *O*-acetyl groups to xylan or mannan oligomers. We developed an *in vitro* assay based on the same principles as the xylosyltransferase assay, using acetyl-coenzyme A (acetyl-CoA) as the acetyl donor and Xyl₆-2AB or Man₅-2AB as the acceptor (Figure 3a). MALDI-TOF MS analysis of the reaction products indicated that this enzyme is able to transfer up to three acetyl groups to Xyl₆-2AB, generating coenzyme A (CoA) as a by-product (Figures 3b and S6b); however, no acetylation of Man₅-2AB was observed under the same reaction conditions (Figure S6c), indicating that the enzyme is a xylan *O*-acetyltransferase. To confirm the structure of the *O*-acetylated xylooligosaccharides, we performed 2D ¹H NMR analysis of the products of scaled-up reactions using unlabeled xylohexaose as the acceptor substrate. The resulting spectra contained signals diagnos-

Figure 3. TBL29/ESK1-catalyzed acetylation of xylooligosaccharides using acetyl-CoA as the donor.

(a) *In vitro* *O*-acetyltransferase reaction scheme. (b) MALDI-TOF MS of the products after incubating TBL29/ESK1 for 16 h with Xyl₆-2AB and acetyl-CoA. Each transfer of an *O*-acetyl (Ac) group increases the mass of the Xyl₆-2AB acceptor by 42 Da to generate products corresponding to the annotated [M + H]⁺ ions. (c) Quantities of acetylated xylosyl residues at O-2 (X2Ac) or O-3 (X3Ac) formed during the first 6 h of the TBL29/ESK-catalyzed reaction. The product concentrations were determined by real-time ¹H NMR analysis.



tic for monosubstituted (2-*O*-acetyl and 3-*O*-acetyl) xylosyl residues, but no signals corresponding to disubstituted xylosyl residues (Figure S7). Thus, the *in vitro* activity of recombinant TBL29/ESK1 is consistent with the reduced xylan acetylation chemotype of *esk1* mutants, which produce xylan with fewer monoacetylated xylosyl residues but no apparent change in disubstituted xylosyl residues (Yuan *et al.*, 2013). To establish the regiospecificity of TBL29/ESK1, we performed real-time NMR analysis of the acetylation reaction. Signals corresponding to monoacetylation at *O*-2 dominated during the initial stages of the reaction, and signals corresponding to monoacetylation at *O*-3 were observed only after a lag period of approximately 30 min (Figures 3c and S8). This is consistent with the enzyme-catalyzed transfer of *O*-acetyl groups to *O*-2, and the spontaneous migration of the acetates between *O*-2 and *O*-3, approaching an equilibrium state in which acetylation at *O*-3 ultimately dominates. Together, our data establish that TBL29/ESK1 catalyzes the transfer of *O*-acetyl groups from acetyl-CoA to *O*-2 of xylosyl residues of β -(1,4) xylooligosaccharides *in vitro*, without the addition of any co-factors or divalent metals (Figure S9). The observation that TBL29/ESK1 is a metal-independent enzyme that can use acetyl-CoA as a donor is comparable with the biochemical analysis of PatB, which has also been shown to be a metal-independent *O*-acetyltransferase capable of using a variety of simple acetyl donors *in vitro* (Moynihan and Clarke, 2014). The availability of the recombinant enzymes and robust biochemical assays reported here will enable research aimed at resolving the complex molecular mechanisms that plants use for polysaccharide *O*-acetylation *in vivo*.

Our data establish the biochemical function of two crucial enzymes in the synthesis and substitution of one of the world's most abundant glycopolymers, xylan. Xylan biosynthesis synthesis, like that of other hemicellulosic polymers, occurs in the Golgi apparatus. Accordingly,

membrane topology studies indicate that the catalytic domain of IRX10-L is localized in the Golgi lumen (Søgaard *et al.*, 2012), and TBL29/ESK1 is predicted to be a type-II membrane protein also localized in the Golgi apparatus (Yuan *et al.*, 2013). This led us to hypothesize that we could synthesize decorated polymers *in vitro* through the sequential or combined catalytic activity of these two proteins. To test this approach we first generated large (up to DP 21) 2AB-labeled xylooligosaccharides *in vitro* using IRX10-L (Figure 4a), then subjected them to TBL29/ESK1-catalyzed acetylation. MALDI-TOF MS analysis of the products demonstrated that TBL29/ESK1 adds up to five acetyl substituents to xylooligosaccharides with DP ≥ 13 (Figure 4b). These findings led us to investigate whether IRX10-L and TBL29/ESK1 may act synergistically *in vitro*, increasing the solubility of nascent polymeric xylan through *O*-acetylation to produce polymers that are larger than those made by IRX10-L alone. Contrary to our expectation, however, the prior introduction of acetyl groups limited the ability of IRX10-L to extend the acceptor *in vitro* (Figure S10). This effect does not necessarily hinder xylan elongation *in vivo*, where, for example, the addition of each xylosyl residue to the polymer may be temporally or spatially isolated from its acetylation.

In summary, we have demonstrated that IRX10-L has UDP-Xyl: β -(1,4)-xylosyl transferase activity, and on this basis we propose the name XYLAN SYNTHASE-1 (XYS1) for this enzyme. Furthermore, we have shown TBL29/ESK1 catalyzes the subsequent addition of *O*-acetyl groups from acetyl-CoA to the 2-position of xylosyl backbone residues, making it the only plant polysaccharide *O*-acetyl transferase that has been biochemically characterized to date. We thus propose the name XYLAN *O*-ACETYLTRANSFERASE 1 (XOAT1) for this enzyme. Our data make it unlikely but do not rule out the possibility (Lee *et al.*, 2012) that the GT43 proteins IRX9 and IRX14 also catalyze the transfer of




Figure 4. MALDI-TOF MS spectra of xylan synthesized and acetylated *in vitro*.

(a) Products formed after incubating IRX10-L for 96 h with Xyl₆-2AB and UDP-Xyl under standard assays conditions. Annotated $[M + H]^+$ ions correspond to xylooligosaccharides with between six and 21 residues.

(b) Products of the reaction described in (a) were incubated with TBL29/ESK1 and acetyl-CoA for 16 h. The number of acetates introduced to each oligosaccharide is indicated by asterisks above its $[M + H]^+$ ion.

xylosyl residues from a donor to the elongating xylan chain. Rather, we propose that these proteins increase xylan synthesis *in vivo*, either by playing a role in chain initiation or by a structural role such as the formation of a multimeric xylan synthase complex. Notably, IRX10 and IRX10-L are both predicted to contain signal peptides rather than transmembrane domains (Wu *et al.*, 2009). Hence, the possibility cannot be discounted that IRX9 and/or IRX14 actually play structural rather than catalytic roles in xylan synthesis, akin to the role that the catalytically inactive, membrane-anchored GT8 protein GAUT7 plays by tethering the α -1,4-galacturonosyltransferase GAUT1 to the Golgi membrane, forming a core complex for pectin biosynthesis (Atmodjo *et al.*, 2011). Notably, Arabidopsis IRX9 and its orthologs in rice and poplar lack a conserved DXD motif (Chiniquy *et al.*, 2013).

Several lines of evidence suggest that there may be variations in the make-up of the xylan synthase complexes depending upon the xylan structures produced in different tissues and species (York and O'Neill, 2008; Faik *et al.*, 2014; Rennie and Scheller, 2014). Our data do not preclude the participation of other enzymes or co-factors that may be required for xylan backbone synthesis *in vivo*; however, the characterization of XYS1 and XOAT1 described here are key advances towards understanding the complex mechanisms used by plants to synthesize xylans and other plant polysaccharides. Our findings also set the stage for the generation of advanced renewable biomaterials developed and produced using natural or engineered catalysts based on these biosynthetic enzymes.

EXPERIMENTAL PROCEDURES

Generation and purification of fusion proteins

Truncated coding-region sequences of AtIRX10-L (amino acids 22–415), AtIRX9 (amino acids 49–351), AtIRX14 (amino acids 89–525) and AtTBL29/AtESK1 (amino acids 57–486) were amplified from cDNA prepared from inflorescence stems of wild-type *A. thaliana* (Col-0) as a template (Urbanowicz *et al.*, 2012). All reactions were carried out using Phusion High-Fidelity DNA Polymerase (Thermo Scientific, <http://www.thermoscientific.com>). To create Gateway entry clones, *attB*-PCR products were generated using two-step adapter PCR, as described in the Gateway Technology Manual (Life Technologies, <http://www.lifetechnologies.com>). The following primer pairs were used for gene-specific amplification of AtIRX10-L, AtIRX9, AtIRX14 and AtTBL29/AtESK1 sequences: IRX10L_pDONR_GS-F, 5'-AACTTGACTTTCAAGGCTTCCGTCTGAGTCCGGAGTC-3'; IRX10L_pDONR_GS-R, 5'-ACAAGAAAGCTGGGTCTACCAAGGTTTCAGATCAGC-3'; IRX14_pDONR_GS-F, 5'-AACTTGACTTTCAAGGCGATGTTACACAATCCGG-3'; IRX14_pDONR_GS-R, 5'-ACAAGAAAGCTGGGTCTACCAAGGTTTCAGATCAGC-3'; IRX9_pDONR_GS-F, 5'-AACTTGACTTTCAAGGCTTATACTTCTACAAAATC-3'; IRX9_pDONR_GS-R, 5'-ACAAGAAAGCTGGGTCTAGGTGCTTAAACGTGTTCTTG-3'; ESK1_pDONR_GS-F, 5'-AACTTGACTTTCAAGGCTCAAACTCACGACGTC-3'; ESK1_pDONR_GS-R, 5'-ACAAGAAAGCTGGGTCTAACGGGAAATGATACGTGT-3'. Underlined sequences denote the partial *attB* adapter sequences appended to

primers used in the first round of PCR amplification. A second set of universal primers, *attB*_Adapter-F, 5'-GGGGACAAGTTTGTACAAAAAGCAGGCTCTGAAAACCTGTACTTTCAAGGC-3', and *attB*_Adapter-R, 5'-GGGGACCCTTTGTACAAGAAAGCTGGGTC-3', was used to insert a tobacco etch virus (TEV) protease cleavage site and complete the *attB* recombination region. The completed *attB*-PCR products were gel purified using the AxyPrep DNA Gel Extraction Kit (Corning, <http://www.corning.com>) and cloned into the pDONR221 plasmid vector (Life Technologies) using Gateway BP Clonase II Enzyme Mix (Life Technologies), following the manufacturer's instructions. The transformants were confirmed by sequence analysis using M13 universal primers M13 Forward (-20) and M13 Reverse. To generate expression clones, the entry clones were recombined into a Gateway-adapted version of the pGen2 mammalian expression vector (pGen2-DEST; Meng *et al.*, 2013), using Gateway LR Clonase II Enzyme Mix (Life Technologies), and confirmed by sequence analysis. The resulting expression construct encodes a fusion protein comprised of an NH₂-terminal signal sequence, an 8xHis tag, an AviTag recognition site, the 'superfolder' GFP coding region and the 7-amino acid recognition sequence of the TEV protease, followed by the corresponding enzyme catalytic domain. For transfection, plasmids were purified using the PureLink HiPure Plasmid Filter Maxiprep Kit (Life Technologies).

Recombinant enzyme expression was accomplished by transient transfection of HEK293 cells, as previously described (Meng *et al.*, 2013). Briefly, suspension culture HEK293f cells (FreeStyle 293-F cells; Life Technologies) were maintained in serum-free Freestyle 293 expression medium (Life Technologies). Transfections were initiated at cell densities of 2.5×10^6 by the addition of $3 \mu\text{g ml}^{-1}$ of the respective expression plasmid and $9 \mu\text{g ml}^{-1}$ polyethylenimine to the culture (linear 25-kDa PEI; Polysciences, <http://www.polysciences.com>). After 24 h the cultures were diluted 1:1 with Freestyle 293 expression medium containing valproic acid (2.2 mM final concentration), and protein production was continued for a further 4–5 days at 37°C.

All chromatography experiments were carried out on an ÄKTA FPLC system (GE Healthcare, <http://www.gehealthcare.com>). Purification of His8-GFP-tagged enzymes secreted into the culture medium by HEK293 cells was performed using His TALON Superflow Cartridges (Clontech, <http://www.clontech.com>) or HisTrap HP columns (GE Healthcare), according to the manufacturer's instructions. The purity of the proteins was confirmed by SDS-PAGE, Coomassie Brilliant Blue R-250 (Bio-Rad, <http://www.bio-rad.com>) staining and in-gel GFP fluorescence. For activity assays, proteins were buffer-exchanged into HEPES sodium salt-HCl (75 mM), pH 6.8, using either a PD-10 gel filtration column (GE Healthcare) or dialysis (3500 molecular weight cut-off), and then concentrated using a 30-kDa molecular weight cut-off Amicon Ultra centrifugal filter device (Millipore, now Merck Millipore, <http://www.merckmillipore.com>). To examine the divalent metal dependence of IRX10-L and TBL29/ESK1, metal-depleted enzyme was prepared by dialysis against HEPES sodium salt-HCl (75 mM) buffer, pH 6.8, containing 5 g L^{-1} Chelex-100 (Bio-Rad) at 4°C for a minimum of 6 h. For real-time NMR experiments, purified proteins were buffer-exchanged into potassium bicarbonate (50 mM), pH 6.8, prepared with D₂O (99.98%; Cambridge Isotope Laboratories, <http://www.isotope.com>) by diafiltration using a 30-kDa molecular weight cut-off Amicon Ultra centrifugal filter device (Millipore). Protein concentration was determined by the Bio-Rad protein assay (Bio-Rad).

Determination of xylosyltransferase activity

The 1,4- β -D-xylooligosaccharides, 1,4- β -D-mannooligosaccharides, 1,4- β -D-glucosaccharides and the endo-1,4- β -xylanase M1

(*Trichoderma viride*) were obtained from Megazyme International (<http://www.megazyme.com>). UDP-xylose, UDP-arabinopyranose and UDP-galacturonic acid were purchased from CarboSource Services (<http://www.csrc.uga.edu>), and UDP-galactose, UDP-glucose, UDP-glucuronic acid and acetyl-CoA were purchased from Sigma-Aldrich (<http://www.sigmaaldrich.com>). β -(1,4)-xylohexaose and β -(1,4)-mannopentaose were labeled at the reducing terminus with 2-aminobenzamide (2AB), as described previously (Ishii *et al.*, 2002). Xylosyltransferase activity was routinely assayed by using an activated sugar donor (UDP-Xyl) and the chemically labeled acceptor 2AB- β -(1,4)-xylohexaose (Xyl₆-2AB). The standard xylosyltransferase assay (15 μ l) was performed in HEPES sodium salt-HCl (75 mM) buffer, pH 6.8, containing 3.8 μ g of purified protein (IRX10-L, IRX9, IRX14), UDP-Xyl (2 mM) as donor and Xyl₆-2AB (0.25 mM) as acceptor. The standard assays were performed in the presence of MnCl₂ (1 mM) and MgCl₂ (1 mM), or only MgCl₂ (1 mM). Reactions were incubated at 25°C for the indicated times. For the determination of the optimal pH, standard reactions were carried out at pH 5.5–6.7 in 2-(*N*-morpholine)-ethanesulphonic acid (MES)-KOH (50 mM) and at pH 6.8–8.0 in HEPES buffer (50 mM). To determine the effect of divalent cations on IRX10-L activity, metal-depleted enzyme and donor substrate were incubated for 30 min at 25°C in the presence of EDTA, MgCl₂, MnCl₂, CaCl₂, CoCl₂ (5 mM) or without additives. The reactions were then initiated by the addition of the acceptor substrate (Xyl₆-2AB) and allowed to proceed for 16 h. To determine whether IRX10-L can use other UDP sugars as sugar donors, IRX10-L and Xyl₆-2AB (0.25 mM) were incubated for 16 h in the presence of each of the following activated nucleotide sugars: 2 mM of UDP-arabinopyranose, UDP-galactose, UDP-galacturonic acid, UDP-glucose or UDP-glucuronic acid. Reaction products were analyzed by MALDI-TOF MS and quantified by RP-HPLC, as described below.

MALDI-TOF MS analysis

The reaction products were analyzed by MALDI-TOF MS using an LT Bruker LT Microflex spectrometer (<http://www.bruker.com>). Aliquots (5 μ l) of the reaction mixture were incubated with 1 μ l of a suspension of Dowex-50 cation exchanger resin in water for 1 h. After centrifugation, 1 μ l of the supernatants was mixed with an equal volume of matrix solution (20 mg ml⁻¹ 2,5-dihydroxybenzoic acid in aqueous 50% methanol) on the target plate. The positive-ion spectra were recorded and at least 200 laser shots were summed to generate each spectrum.

RP-HPLC analysis and quantification

The xylosyltransferase activity was measured using RP-HPLC analysis of the fluorescent 2AB-labeled oligosaccharides generated by IRX10-L. HPLC was performed with an Agilent 1100 series LC system (<http://www.agilent.com>) and a Shimadzu RF-10AXL fluorescence detector (<http://www.shimadzu.com/>). The reaction products were separated using a Luna C18 column (250-mm long, 4.6-mm inner diameter). The column was eluted at 0.5 ml min⁻¹ with aqueous 20% methanol for 30 min followed by a 5-min gradient to 80% methanol, which was held for 5 min. The column was re-equilibrated by a 5-min gradient to return to the initial conditions and washed for an additional 5 min in these conditions. The elution of fluorescent oligosaccharides was monitored using a fluorescence detector (330-nm excitation and 420-nm emission). The molar concentrations of products were quantified assuming that the molar fluorescence response was the same for all labeled substrates and products, and that the total molar concentration of these labeled molecules was equal to the initial concentration of the Xyl₆-2AB acceptor. Each detected oligosaccharide was assigned an index i , which

corresponds to the DP of the oligosaccharide. The signal s_i of each oligosaccharide peak was integrated and used to calculate molar concentrations c_i by the following equation: $c_i = \frac{s_i}{S} [\text{acceptor}]$, where $S = \sum_{i=6}^{i=18} S_i$ and [acceptor] is the initial molar concentration of the Xyl₆-2AB acceptor. The molar quantity a_i (nmol) of each product produced per microgram of protein was calculated by dividing its concentration by the concentration of protein (0.25 μ g μ l⁻¹) in the reaction $a_i = \frac{c_i}{[\text{enzyme}]}$, where [enzyme] is the concentration of the enzyme. The total quantity x of xylosyl residues (nmol) transferred per microgram of protein was quantified by multiplying the molar quantity a_i of each product per microgram of protein by its extension number $n_i = i - 6$, where 6 is the DP of the acceptor substrate, and summing the result: $x = \sum_{i=6}^{i=18} n_i a_i$. Theoretical product population calculations were also performed for each time point. First, the observed abundances a_i of the oligosaccharides were converted to fractional populations $f_i = \frac{a_i}{A}$, where $A = \sum_{i=6}^{i=18} a_i$. The Poisson expectation value λ was calculated by multiplying each fractional population by the extension number n_i and summing the result: $\lambda = \sum_{i=6}^{i=18} n_i f_i$. The value of λ corresponds to the average number of xylosyl residues transferred per oligosaccharide acceptor molecule in the reaction. The theoretical fractional population of each oligosaccharide p_i was calculated as a Poisson probability $P(n_i|\lambda)$ for each oligosaccharide in the standard way: $p_i = P(n_i|\lambda) = \frac{\lambda^{n_i} e^{-\lambda}}{n_i!}$. The observed and theoretical fractional populations for the products of the reaction catalyzed by IRX10-L were remarkably similar.

Structural characterization of xylooligosaccharides synthesized *in vitro*

NMR spectroscopy was used to structurally characterize the *in vitro* synthesized β -(1,4)-xylooligosaccharides. Scaled-up enzyme-catalyzed reactions were carried out in potassium bicarbonate (50 mM), pH 6.8 in D₂O containing unlabeled xylohexaose (100 μ g) and UDP-Xyl (2 mM) as the donor. The course of the reaction was monitored by real-time 1D ¹H NMR. After 120 h, the products of the scaled-up reaction were fractionated by SEC on a Superdex-75 HR10/30 SEC column (GE Healthcare) eluted with water. The fractions containing the xylo-oligosaccharides were lyophilized, dissolved in D₂O (0.3 ml, 99.9%; Cambridge Isotope Laboratories, <http://www.isotope.com>) and analyzed by 1D and 2D NMR spectroscopy. Data were recorded at 298 K with an Agilent-NMR spectrometer operating at 600 MHz, equipped with a 5-mm NMR cold probe. The 2D homonuclear and heteronuclear experiments were recorded using standard Varian pulse programs. Chemical shifts were measured relative to internal acetone (δ ¹H 2.225; δ ¹³C 30.89). Data were processed using MESTReNOVA (Universidad de Santiago de Compostela, <http://www.usc.es>). Long-range scalar couplings in the gHMBC spectrum were used to establish the glycosidic linkage positions between the xylosyl residues in the reaction products.

The products of the xylosyltransferase reaction catalyzed by IRX10-L for 16 h, using Xyl₆-2AB as the acceptor, and under the same conditions described for the standard xylosyltransferase assay, were treated for 1 h with the endo-1,4- β -xylanase M1. Aliquots of the reaction with and without the endoxylanase treatment were analyzed by RP-HPLC, as described above.

Kinetics analysis and determination of the acceptor DP preferences of IRX10-L

The quantity of UDP formed as a by-product of the xylosyltransferase reactions was determined using the UDP-Glo™ Glycosyltransferase Assay (Promega, <http://www.promega.com>), according to the manufacturer's instructions. For the determination of kinetic

parameters, xylosyltransferase reactions (5 μ l) comprising HEPES sodium salt-HCl (75 mM) buffer, pH 6.8, and purified IRX10-L (1.25 μ g) were carried out using UDP-xylose (800 μ M; CarboSource Services) as a donor substrate and several different concentrations of xylohexaose (0–4.0 mM) as an acceptor. Reactions were performed at 26°C for up to 120 min, and were stopped at the indicated times by flash-freezing in liquid nitrogen. For the determination of protein concentration dependence, xylosyltransferase reactions were carried out with UDP-xylose (800 μ M) as a donor, xylohexaose (1.0 mM) as an acceptor and increasing concentrations of purified IRX10-L (0–1.25 μ g) for 60 min at 26°C. For the detection of UDP, reactions were mixed with an equal volume of UDP-Glo Detection Reagent (5 μ l) in white polystyrene, low-volume, 384-well assay plates (Corning, <http://www.corning.com>), and incubated for 60 min at 23°C. After incubation, luminescence measurements were performed using a multifunctional microplate reader (POLARstar OPTIMA; BMG Labtech, <http://www.bmg-labtech.com>). A standard curve was used for the quantification of UDP produced, and the detection limit of the luminescent assay was 40 nM UDP with a linear response up to 20 μ M. The steady-state parameters K_m and V_{max} were calculated by fitting the initial velocities with the Michaelis–Menten equation using nonlinear curve fitting in PRISM 6 (GraphPad, <http://www.graphpad.com>).

To investigate xylosyltransferase activity on xylooligomer acceptors of different degrees of polymerization, reactions were performed with the following unlabeled acceptor substrates: xylose (Xyl₁), xylobiose (Xyl₂), xylotriose (Xyl₃), xylo-tetraose (Xyl₄), xylopentaose (Xyl₅), xylohexaose (Xyl₆) and cellopentaose (Glc₅). Each reaction (5 μ l) was carried out for 12 h at 25°C and contained 0.4 μ g of enzyme, UDP-Xyl (100 μ M) donor and the indicated acceptor substrate (65 μ M). Glycosyltransferase activity was measured using the bioluminescent UDP-Glo™ Glycosyltransferase Assay (Promega), as described above.

Determination of xylan O-acetyltransferase activity

Xylan O-acetyltransferase activity was routinely assayed by using acetyl-CoA as a donor and 2AB-Xyl₆ or xylohexaose as acceptor substrates. The standard assay (15 μ l) was performed in HEPES sodium salt-HCl (75 mM) buffer, pH 6.8, containing purified protein (4 μ M), acetyl-CoA (1 mM) donor and Xyl₆-2AB (0.25 mM) acceptor, unless indicated. Mannan O-acetyltransferase activity was assayed as described above using 2AB- β -(1,4)-mannopentaose (Man₅-2AB) as an acceptor. The reaction products were analyzed by MALDI-TOF MS, as described above. The optimal acetyl-CoA concentration and the effect of divalent cations on TBL29/ESK1 activity were determined using the same experimental procedures described for IRX10-L. For structural analysis of the reaction products, unlabeled xylohexaose (100 μ g) and acetyl-CoA (2 mM) were incubated with TBL29/ESK1 in 50 mM potassium bicarbonate (50 mM), pH 6.8, in D₂O for 48 h. The high-resolution gCOSY was recorded as previously described (Jia *et al.*, 2005). Resonances were assigned to glycosyl residues based on our analysis of acetylated xylooligosaccharides released from Arabidopsis stem-cell walls by the endo-1,4- β -xylanase M1 and on published data (Teleman *et al.*, 2000; Kabel *et al.*, 2003).

To investigate the positional specificity of TBL29/ESK1, the reaction catalyzed by TBL29/ESK1 *in vitro* was monitored using real-time ¹H NMR. The assay was performed at 25°C in potassium bicarbonate (50 mM), pH 6.8, in D₂O containing unlabeled xylohexaose (100 μ g), acetyl-CoA (2 mM) and 50 μ g of purified TBL29/ESK1. Data acquisition was initiated 5 min after the reaction components were mixed. 1D ¹H spectra consisted of 64 transients that were acquired with water pre-saturation every

10 min over a 12-h time period, generating a spectral array representing the progress of the reaction. The resonance peaks corresponding to the methyl protons of the acetyl groups attached to O-2 or O-3 of internal xylosyl residues were identified in the arrayed spectra, and their areas were determined by integration or by deconvolution in the case of overlapping peaks. The concentrations of the acetylated products were calculated based on the initial concentration of acetyl-CoA added to the reaction. After monitoring the reaction for 12 h, the products were analyzed by 1D and 2D NMR, as described above. All data processing was performed using MESTRENOVA (Universidad de Santiago de Compostela).

ACKNOWLEDGEMENTS

We thank M.A. O'Neill, Peter J. Smith and J. Backe for technical support, and the Center for Plant and Microbial Complex Carbohydrates (DEFG0296ER20097) for equipment support. B.R.U., M.J.P. and W.S.Y. were funded by The BioEnergy Science Center (BESC), a US Department of Energy Bioenergy Research Center supported by the Office of Biological and Environmental Research in the U.S. Department of Energy Office of Science. H.A.M. and K.W.M. were supported by the National Institutes of Health (grant P41GM103390). The nucleotide sugars obtained from CarboSource Services, which is supported in part by the DOE-funded Center For Plant and Microbial Complex Carbohydrates (grant DE-FG02-93ER20097). All authors contributed to the intellectual development and/or writing of the article. All authors designed the research, and B.R.U., M.J.P., and H.A.M. performed the experiments.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

Figure S1. Characteristics of the purified recombinant IRX10-L.

Figure S2. Xylosyltransferase activity assays of several enzymes predicted to be involved in xylan synthesis.

Figure S3. Structural characterization of the products of the reaction catalyzed by IRX10-L.

Figure S4. Heteronuclear ¹H-¹³C NMR characterization of the products generated by IRX10-L.

Figure S5. Determination of the optimal pH for xylosyltransferase activity.

Figure S6. Substrate specificity of the purified recombinant TBL29/ESK1.

Figure S7. Characterization by NMR of the products of the reaction catalyzed by TBL29/ESK1.

Figure S8. Determination of the regio-specificity of TBL29/ESK1.

Figure S9. Effect of divalent metals on the acetyltransferase activity of TBL29/ESK1.

Figure S10. Effect of acceptor substrate O-acetylation on the xylosyltransferase activity of IRX10-L.

REFERENCES

- Atmodjo, M.A., Sakuragi, Y., Zhu, X., Burrell, A.J., Mohanty, S.S., Atwood, J.A. III, Orlando, R., Scheller, H.V. and Mohnen, D. (2011) Galacturonosyltransferase (GAUT)1 and GAUT7 are the core of a plant cell wall pectin biosynthetic homogalacturonan: galacturonosyltransferase complex. *Proc. Natl Acad. Sci. USA*, **108**, 20225–20230.
- Brown, D.M., Zeef, L.A.H., Ellis, J., Goodacre, R. and Turner, S.R. (2005) Identification of novel genes in Arabidopsis involved in secondary cell wall formation using expression profiling and reverse genetics. *Plant Cell*, **17**, 2281–2295.

- Brown, D.M., Zhang, Z., Stephens, E., Dupree, P. and Turner, S.R. (2009) Characterization of IRX10 and IRX10-like reveals an essential role in glucuronoxylan biosynthesis in Arabidopsis. *Plant J.* **57**, 732–746.
- Burton, R.A., Wilson, S.M., Hrmova, M., Harvey, A.J., Shirley, N.J., Stone, B.A., Newbigin, E.J., Bacic, A. and Fincher, G.B. (2006) Cellulose synthase-like CslF genes mediate the synthesis of cell wall (1,3;1,4)-beta-D-glucans. *Science*, **311**, 1940–1942.
- Chiniquy, D., Varanasi, P., Oh, T. et al. (2013) Three novel rice genes closely related to the Arabidopsis IRX9, IRX9L, and IRX14 genes and their roles in xylan biosynthesis. *Front. Plant Sci.* **4**, 83.
- Cocuron, J.-C., Lerouxel, O., Drakakaki, G., Alonso, A.P., Liepman, A.H., Keegstra, K., Raikhel, N. and Wilkerson, C.G. (2007) A gene from the cellulose synthase-like C family encodes a beta-1,4 glucan synthase. *Proc. Natl Acad. Sci. USA*, **104**, 8550–8555.
- Dhugga, K.S., Barreiro, R., Whitten, B. et al. (2004) Guar seed beta-mannan synthase is a member of the cellulose synthase super gene family. *Science*, **303**, 363–366.
- Doblin, M.S., Pettolino, F.A., Wilson, S.M., Campbell, R., Burton, R.A., Fincher, G.B., Newbigin, E. and Bacic, A. (2009) A barley cellulose synthase-like CSLH gene mediates (1,3;1,4)-beta-D-glucan synthesis in transgenic Arabidopsis. *Proc. Natl Acad. Sci. USA*, **106**, 5996–6001.
- Faik, A., Jiang, N. and Held, M. (2014) Xylan Biosynthesis in Plants, Simply Complex. In *Plants and BioEnergy* (McCann, M.C., Buckeridge, M.S. and Carpita, N.C., eds). New York, NY: Springer, pp. 153–181.
- Geshi, N., Harholt, J., Sakuragi, Y., Jensen, J.K. and Scheller, H.V. (2010) Glycosyltransferases of the GT47 family. In *Annual Plant Reviews: Plant Polysaccharides, Biosynthesis and Bioengineering*. vol. 41, (Ulvskov, P., ed). West Sussex: Wiley-Blackwell, pp. 265–283.
- Gille, S. and Pauly, M. (2012) O-acetylation of plant cell wall polysaccharides. *Front. Plant Sci.* **3**, 12.
- Gille, S., de Souza, A., Xiong, G., Benz, M., Cheng, K., Schultink, A., Reza, I.-B. and Pauly, M. (2011) O-acetylation of Arabidopsis hemicellulose xyloglucan requires AXY4 or AXY4L, proteins with a TBL and DUF231 domain. *Plant Cell*, **23**, 4041–4053.
- Ishii, T., Ichita, J., Matsue, H., Ono, H. and Maeda, I. (2002) Fluorescent labeling of pectic oligosaccharides with 2-aminobenzamide and enzyme assay for pectin. *Carbohydr. Res.* **337**, 1023–1032.
- Jensen, J.K., Kim, H., Cocuron, J.-C., Orlor, R., Ralph, J. and Wilkerson, C.G. (2011) The DUF579 domain containing proteins IRX15 and IRX15-L affect xylan synthesis in Arabidopsis. *Plant J.* **66**, 387–400.
- Jensen, J.K., Johnson, N. and Wilkerson, C.G. (2013) Discovery of diversity in xylan biosynthetic genes by transcriptional profiling of a heteroxylan containing mucilaginous tissue. *Front. Plant Sci.* **4**, 183.
- Jia, Z., Cash, M., Darvill, A.G. and York, W.S. (2005) NMR characterization of endogenously O-acetylated oligosaccharides isolated from tomato (*Lycopersicon esculentum*) xyloglucan. *Carbohydr. Res.* **340**, 1818–1825.
- Kabel, M.A., de Waard, P., Schols, H.A. and Voragen, A.G.J. (2003) Location of O-acetyl substituents in xylo-oligosaccharides obtained from hydrothermally treated Eucalyptus wood. *Carbohydr. Res.* **338**, 69–77.
- Kuroyama, H. and Tsumuraya, Y. (2001) A xylosyltransferase that synthesizes beta-(1 → 4)-xylans in wheat (*Triticum aestivum* L.) seedlings. *Planta*, **213**, 231–240.
- Lairson, L.L., Henrissat, B., Davies, G.J. and Withers, S.G. (2008) Glycosyltransferases: structures, functions, and mechanisms. *Annu. Rev. Biochem.* **77**, 521–555.
- Lee, C., Zhong, R. and Ye, Z.-H. (2012) Arabidopsis family GT43 members are xylan xylosyltransferases required for the elongation of the xylan backbone. *Plant Cell Physiol.* **53**, 135–143.
- Liu, J. and Mushegian, A. (2003) Three monophyletic superfamilies account for the majority of the known glycosyltransferases. *Protein Sci.* **12**, 1418–1431.
- Liwanag, A.J.M., Ebert, B., Verherbruggen, Y., Rennie, E.A., Rautengarten, C., Oikawa, A., Andersen, M.C.F., Clausen, M.H. and Scheller, H.V. (2012) Pectin biosynthesis: GAL51 in Arabidopsis thaliana is a beta-1,4-galactan beta-1,4-galactosyltransferase. *Plant Cell*, **24**, 5024–5036.
- Manabe, Y., Verherbruggen, Y., Gille, S. et al. (2013) Reduced wall acetylation proteins play vital and distinct roles in cell wall O-acetylation in Arabidopsis. *Plant Physiol.* **163**, 1107–1117.
- May, J.F., Splain, R.A., Brotschi, C. and Kiessling, L.L. (2009) A tethering mechanism for length control in a processive carbohydrate polymerization. *Proc. Natl Acad. Sci. USA*, **106**, 11851–11856.
- Meng, L., Forouhar, F., Thieker, D. et al. (2013) Enzymatic basis for N-glycan sialylation: structure of rat alpha2,6-sialyltransferase (ST6GAL1) reveals conserved and unique features for glycan sialylation. *J. Biol. Chem.* **288**, 34680–34698.
- Moynihan, P.J. and Clarke, A.J. (2010) O-acetylation of peptidoglycan in gram-negative bacteria identification and characterization of peptidoglycan O-acetyltransferase in *Neisseria gonorrhoeae*. *J. Biol. Chem.* **285**, 13264–13273.
- Moynihan, P.J. and Clarke, A.J. (2013) Assay for peptidoglycan O-acetyltransferase: a potential new antibacterial target. *Anal. Biochem.* **439**, 73–79.
- Moynihan, P.J. and Clarke, A.J. (2014) Substrate specificity and kinetic characterization of peptidoglycan O-acetyltransferase B from *Neisseria gonorrhoeae*. *J. Biol. Chem.* **289**, 16748–16760.
- Pear, J.R., Kawagoe, Y., Schreckengost, W.E., Delmer, D.P. and Stalker, D.M. (1996) Higher plants contain homologs of the bacterial celA genes encoding the catalytic subunit of cellulose synthase. *Proc. Natl Acad. Sci. USA*, **93**, 12637–12642.
- Pena, M.J., Zhong, R., Zhou, G.-K., Richardson, E.A., O'Neill, M.A., Darvill, A.G., York, W.S. and Ye, Z.-H. (2007) Arabidopsis irregular xylem8 and irregular xylem9: implications for the complexity of glucuronoxylan biosynthesis. *Plant Cell*, **19**, 549–563.
- Porchia, A.C. and Scheller, H.V. (2000) Arabinoxylan biosynthesis: identification and partial characterization of beta-1,4-xylosyltransferase from wheat. *Physiol. Plant.* **110**, 350–356.
- Rennie, E.A. and Scheller, H.V. (2014) Xylan biosynthesis. *Curr. Opin. Biotechnol.* **26**, 100–107.
- Rennie, E.A., Hansen, S.F., Baidoo, E.E.K., Hadi, M.Z., Keasling, J.D. and Scheller, H.V. (2012) Three members of the Arabidopsis glycosyltransferase family 8 are xylan glucuronosyltransferases. *Plant Physiol.* **159**, 1408–1417.
- Richmond, T.A. and Somerville, C.R. (2000) The cellulose synthase superfamily. *Plant Physiol.* **124**, 495–498.
- Sandhu, A.P.S., Randhawa, G.S. and Dhugga, K.S. (2009) Plant cell wall matrix polysaccharide biosynthesis. *Mol. Plant.* **2**, 840–850.
- Scheller, H.V. and Ulvskov, P. (2010) Hemicelluloses. *Annu. Rev. Plant Biol.* **61**, 263–289.
- Simkovic, I. (2013) Unexplored possibilities of all-polysaccharide composites. *Carbohydr. Polym.* **95**, 697–715.
- Søgaard, C., Stenbaek, A., Bernard, S., Hadi, M., Driouch, A., Scheller, H.V. and Sakuragi, Y. (2012) GO-PROMTO illuminates protein membrane topologies of glycan biosynthetic enzymes in the golgi apparatus of living tissues. *PLoS ONE*, **7**, e31324.
- Teleman, A., Lundqvist, J., Tjerneld, F., Stalbrand, H. and Dahlman, O. (2000) Characterization of acetylated 4-O-methylglucuronoxylan isolated from aspen employing 1H and 13C NMR spectroscopy. *Carbohydr. Res.* **329**, 807–815.
- Urbanowicz, B.R., Pena, M.J., Ratnaparkhe, S. et al. (2012) 4-O-methylation of glucuronic acid in Arabidopsis glucuronoxylan is catalyzed by a domain of unknown function family 579 protein. *Proc. Natl Acad. Sci. USA*, **109**, 14253–14258.
- Wu, A.-M., Rihouey, C., Seveno, M., Hornblad, E., Singh, S.K., Matsunaga, T., Ishii, T., Lerouge, P. and Marchant, A. (2009) The Arabidopsis IRX10 and IRX10-LIKE glycosyltransferases are critical for glucuronoxylan biosynthesis during secondary cell wall formation. *Plant J.* **57**, 718–731.
- Wu, A.-M., Hornblad, E., Voxeur, A., Gerber, L., Rihouey, C., Lerouge, P. and Marchant, A. (2010) Analysis of the Arabidopsis IRX9/IRX9-L and IRX14/IRX14-L pairs of glycosyltransferase genes reveals critical contributions to biosynthesis of the hemicellulose glucuronoxylan. *Plant Physiol.* **153**, 542–554.
- Xiong, G., Cheng, K. and Pauly, M. (2013) Xylan O-Acetylation impacts xylem development and enzymatic recalcitrance as indicated by the Arabidopsis mutant tbl29. *Mol. Plant.* **6**, 1373–1375.
- York, W.S. and O'Neill, M.A. (2008) Biochemical control of xylan biosynthesis – which end is up? *Curr. Opin. Plant Biol.* **11**, 258–265.
- Yuan, Y., Teng, Q., Zhong, R. and Ye, Z.-H. (2013) The Arabidopsis DUF231 domain-containing protein ESK1 mediates 2-O- and 3-O-acetylation of xylosyl residues in xylan. *Plant Cell Physiol.* **54**, 1186–1199.